

THE AMALGAMATED FREE PRODUCT OF SEMIFINITE HYPERFINITE VON NEUMANN ALGEBRAS OVER ATOMIC TYPE I SUBALGEBRAS

DANIEL REDELMEIER

ABSTRACT. In this paper we describe the amalgamated free product of finite and semifinite hyperfinite von Neumann algebras over atomic type I subalgebras. To do this we extend the notions of free dimension and standard embeddings used in the related results for finite von Neumann algebras to ones which work better for the semifinite case. We also define classes $\mathcal{R}_3$ (of finite von Neumann algebras) and $\mathcal{R}_4$ (of semifinite von Neumann algebras) which are closed under such amalgamated free products.

1. INTRODUCTION

Voiculescu's amalgamated free product for C^* and von Neumann algebras in [11] and [10] have proved to be very useful constructions. Early uses include Popa's use of the the amalgamated free product of von Neumann algebras in [9] to construct subfactors with arbitrary allowable index.

The standard free product of finite hyperfinite von Neumann algebras was described by Dykema in [2]. He also described the amalgamated free product of multimatrix algebras in [4]. In [6] we described the amalgamated free product of finite hyperfinite von Neumann algebras over finite dimensional subspaces. Here we will extend this to semifinite hyperfinite von Neumann algebras over atomic type I subalgebras, for which the induced trace is also semifinite.

In [5], Dykema also showed that a certain class of von Neumann algebras (referred to as $\mathcal{R}$ in that paper and $\mathcal{R}_1$ here) was closed under amalgamated free products over finite dimensional subspaces. This was used by Kodiyalam and Sunder in [8] and in the related paper by Guionnet, Jones, and Shlyakhtenko in [7]. This class was extended to a more natural class (which we refer to as $\mathcal{R}_2$) in [6]. Here we extend these to classes $\mathcal{R}_3$ of finite von Neumann algebras and $\mathcal{R}_4$ of semifinite von Neumann algebras which are each closed under amalgamated free products over type I atomic subalgebras.

Key words and phrases. amalgamated free product, hyperfinite von Neumann algebras.

2. BASIC THEOREMS AND DEFINITIONS

2.1. Preliminaries. Throughout this paper we will use finite and semifinite von Neumann algebras, with specified normal faithful semifinite trace. In most cases we will also have a specified atomic type I subalgebra D of these von Neumann algebras. We will in general assume that there is a trace preserving conditional expectation from our main algebras onto D and that the trace will still be semifinite (or finite) on D . If we refer to an expectation without specifying which one, this is the one we mean. Unless specified otherwise we will also be assuming homomorphisms are trace preserving.

With these von Neumann algebras in mind, we will be using the amalgamated free product for von Neumann algebras. Note it is clear from the definition of the trace on $A *_D B$, $\tau_D \circ E_D$, that if the trace on A , B and D is semifinite then it will be on $A *_D B$.

We will often use the following notation $A_1^{p_1} \oplus A_2^{p_2} \oplus \dots$, where the p_i denote the central support of the A_i (i.e. the identity in A_i as part of the larger algebra, also sometimes referred to as the matrix unit). In the case of atomic type I factors, we may also use the notation A_t , where t denotes the trace of a minimal projection in this factor (clearly this doesn't exist in diffuse algebras, and is not necessarily unique if it is not a factor).

Definition 2.1. We will be dealing primarily with four classes of von Neumann algebras, defined as follows:

- Let $\mathcal{R}_1$ (originally $\mathcal{R}$ in [5]) be the class of finite von Neumann algebras which are the finite direct sum of the following types of algebras:
 - (1) Matrix Algebras
 - (2) $M_n \otimes L^\infty([0, 1])$
 - (3) Hyperfinite II_1 factors
 - (4) Interpolated Free Group Factors
- Let $\mathcal{R}_2$ be the class of finite von Neumann algebras which are the direct sum of a finite hyperfinite von Neumann algebra and a finite number of interpolated free group factors (as defined in [6]).
- Let $\mathcal{R}_3$ be the class of finite von Neumann algebras which are the direct sum of a finite hyperfinite von Neumann algebra and a countable number of interpolated free group factors.
- Let $\mathcal{R}_4$ be the class of semifinite von Neumann algebras which are countable direct sums of the following types of algebras:
 - (1) Semifinite hyperfinite von Neumann algebras
 - (2) Interpolated free group factors
 - (3) $F \otimes B(\mathcal{H})$ where F is an interpolated free group factor and $\mathcal{H}$ is a separable Hilbert space.

Note each class strictly contains the previous.

On the class $\mathcal{R}_3$, Dykema defined the notion of *Free Dimension* in [2] in the following way.

Definition 2.2. Let A be a finite von Neumann algebra with specified normal faithful tracial state which is of the format

$$A = H \oplus \bigoplus_{i \in I} L(F_{r_i})^{p_i} \oplus \bigoplus_{j \in J} M_{n_j}^{t_j},$$

where H is a diffuse hyperfinite algebra, the $L(F_{r_i})$ are interpolated free group factors and the M_{n_j} are matrix algebras. The *free dimension* of A (denote $\text{fdim}(A)$) is equal to

$$1 + \left(\sum_{i \in I} \tau(p_i)^2 (r_i - 1) \right) - \sum_{j \in J} t_j^2.$$

This may not be well defined if the interpolated free group factors turn out to be all isomorphic. The definition can be made rigourous by defining it on a generating set, and this is what we mean when we use it. This is not as big a problem as it seems, as in most cases we use the free dimension to determine which interpolated free groups we have, and this would then not matter.

The *standard embedding* was introduced by Dykema in [2].

Definition 2.3. Let $A = L(F_r)$ and $B = L(F_{r'})$ for $r < r'$ and let $\phi : A \rightarrow B$ be a unital embedding. We call ϕ a *standard embedding* if we can find a semicircular system $\omega = \{X_t\}_{t \in T}$ and a copy of R , the hyperfinite II_1 factor free from ω , and $p_t \in R$ so that B is generated by R and $\{p_t X_t p_t\}_{t \in T}$, and such that if $T' = \{t \in T \mid p_t X_t p_t \in \phi(A)\}$ then $\phi(A) = vN(R \cup \{p_t X_t p_t\}_{t \in T'})$.

Standard embeddings have the following properties, proved by Dykema in [2]:

- (1) For $A = L(F_s)$ and $B = L(F_{s'})$, $s < s'$, then for $\phi : A \rightarrow B$ and projection $p \in A$, ϕ is standard if and only if $\phi|_{pAp} \rightarrow \phi(p)B\phi(p)$ is standard.
- (2) The inclusion $A \rightarrow A * B$ is standard if A is an interpolated free group factor and B is an interpolated free group factor, $L(\mathbb{Z})$, or a finite dimensional algebra other than $\mathbb{C}$.
- (3) The composition of standard embeddings is standard.
- (4) For $A_n = L(F_{s_n})$, with $s_n < s_{n'}$ if $n < n'$, and $\phi_n : A_n \rightarrow A_{n+1}$ a sequence of standard embeddings, then the inductive limit of the A_n with the inclusions ϕ_n is L_{F_s} where $s = \lim_{n \rightarrow \infty} s_n$.

2.2. Previous Results and Lemmas. Dykema proved the following as Lemma 4.3 in [4].

Lemma 2.4. Let A and B be von Neumann algebras with subalgebra D , and let $\mathcal{M} = A *_D B$. Let p be a central projection in A , and let $\underline{A} = pD \oplus (1-p)A$ and $\underline{M} = \underline{A} *_D B$. Then the central support of p is the same in $\mathcal{M}$ and $\underline{M}$. Furthermore $p\mathcal{M}p = vN(p\underline{M}p \cup pA) = p\underline{M}p *_D pA$.

Dykema also proved the following, as Lemma 4.2 from [4].

Lemma 2.5. *Let $\mathcal{N} = \overset{r_0}{H} \oplus \bigoplus_{i \in I} \overset{r_i}{F_i} \oplus \bigoplus_{j \in J} \overset{q_j}{M_{n_j}}$, where H is a diffuse hyperfinite algebra. Let $p \in \mathcal{N}$ be a projection such that $\tau(p) = \frac{1}{2}$ and so that neither p nor $1 - p$ is minimal and central in $\mathcal{N}$. Let $D = \overset{p}{\mathbb{C}} \oplus \overset{1-p}{\mathbb{C}}$. Define a matrix algebra $M_2(\mathbb{C})$, where $e_{11} = p$ and $e_{22} = 1 - p$, and let $v = e_{12}$. Then*

$$\mathcal{M} = \mathcal{N} *_D M_2(\mathbb{C}) = F \oplus \bigoplus_{k \in K} \overset{q'_k}{M_{n'_k}},$$

where F is either an interpolated free group factor or a diffuse hyperfinite algebra, and

$$K = \left\{ (j, j') \mid j, j' \in J, j \leq p, j' \leq 1 - p, \frac{t_j}{n_j} + \frac{t_{j'}}{n_{j'}} > \frac{1}{2} \right\}.$$

For $k = (j, j') \in K$, $n_k = 2n_j n_{j'}$ and $t'_k = \frac{n'_k}{2} \left(\frac{t_j}{n_j} + \frac{t_{j'}}{n_{j'}} - \frac{1}{2} \right)$. We also know $\text{fdim}(\mathcal{M}) = \text{fdim}(\mathcal{N}) + \frac{1}{4}$, which then determines F if it is an interpolated free group factor (F is only hyperfinite if $\mathcal{N}$ is dimension 4). Furthermore, the inclusion of $L(F_{s_i}) = r_i \mathcal{N} r_i \rightarrow r_i F r_i$ is standard. for all $i \in I$ (noting if I is not empty, then F is an interpolated free group factor).

The amalgamated free product of multimatrix algebras were described by Dykema as Theorem 5.1 in [4].

Theorem 2.6. *Let A and B be multimatrix algebras with subalgebra D . Then $A *_D B$ is in $\mathcal{R}_3$. If D is finite dimensional it is in $\mathcal{R}_2$ and the hyperfinite part is type I. Furthermore $\text{fdim}(A *_D B) = \text{fdim}(A) + \text{fdim}(B) - \text{fdim}(D)$.*

The following results were proved in [6]

Lemma 2.7. *Let $\mathcal{N} = (M_m \oplus M_n \oplus B) *_D C$ and $\mathcal{M} = (M_{n+m} \oplus B) *_D C$, where B, C are semifinite von Neumann algebras and $D = \bigoplus_{i=1}^K \overset{p_i^D}{\mathbb{C}}$ with $K \in \mathbb{N} \cup \{\infty\}$. $\mathcal{N}$ is included in $\mathcal{M}$ by including M_m and M_n as blocks on the diagonal of M_{n+m} , and B and C by the identity. Assume there exists a partial isometry in $\mathcal{N}$ between minimal projections in M_m and M_n (for example if there exists a factor $\mathcal{F}$ with $M_m \oplus M_n \subseteq \mathcal{F} \subseteq \mathcal{N}$). Then for any minimal projection $p \in M_m$ such that $p \leq p_i^D$ for some i , $p\mathcal{N}p * L(\mathbb{Z}) \cong p\mathcal{M}p$.*

Lemma 2.8. *Let $\mathcal{M} = ((M_n \otimes A) \oplus B) *_D C$ and $\mathcal{N} = (M_n \oplus B) *_D C$ for A is a finite von Neumann algebras and B and C are semifinite, and $D = \bigoplus_{i=1}^K \overset{p_i^D}{\mathbb{C}}, k \in \mathbb{N} \cup \{\infty\}$, where C, B, M_n have expectations onto D and where $E_D^{M_n \otimes A} = E_D^{M_n} \otimes \tau_A$. Let p be a minimal projection in M_n , with $p \leq p_i^D$ for some i . Then $p\mathcal{N}p * A \cong p\mathcal{M}p$.*

Theorem 2.9. *Let A and B be hyperfinite von Neumann Algebras, with finite dimensional subalgebra D . Then $A *_D B$ is in $\mathcal{R}_2$. Furthermore $\text{fdim}(A *_D B) = \text{fdim}(A) + \text{fdim}(B) - \text{fdim}(D)$.*

The following was proved for $\mathcal{R}_1$ in [5] and $\mathcal{R}_2$ in [6]:

Theorem 2.10. *Both $\mathcal{R}_1$ and $\mathcal{R}_2$ are closed under amalgamated free products over finite dimensional subalgebras. In these cases $\text{fdim}(A *_D B) = \text{fdim}(A) + \text{fdim}(B) - \text{fdim}(D)$.*

3. HYPERFINITE VON NEUMANN ALGEBRAS

3.1. Regulated Dimension. Unfortunately free dimension is only defined on finite von Neumann algebras, so here we adapt it in a way that generalizes to semifinite algebras.

For algebras in $\mathcal{R}_3$, where free dimension is defined, we can define the *regulated dimension* in a fairly straight forward way, as follows:

$$\text{rdim}(A) = \tau(I_A)^2(\text{fdim}(A) - 1).$$

This means that for an interpolated free group factor $L(F_t)$ endowed with a trace normalized to λ , the regulated dimension is $\lambda^2(t - 1)$, which is the contribution the interpolated free group factor summand gives in the free dimension formula. For a matrix algebra whose minimal projections have trace t , the regulated dimension is $-t^2$. Finally for any diffuse hyperfinite algebra, the regulated dimension is zero. Unlike the free dimension, the regulated dimension is additive over direct sums (where the trace is preserved in the embeddings), and so the regulated dimension of any algebra in $\mathcal{R}_3$ can easily be determined from the above.

Lemma 3.1. *Let A be an algebra in $\mathcal{R}_3$ and let $p \in A$ be a projection in A with full central support. Then $\text{rdim}(A) = \text{rdim}(pAp)$.*

Proof. First consider an interpolated free group factor $A = L(F_t)$, whose identity has trace λ . Then $\text{rdim}(A) = \lambda^2(t - 1)$. Let $\gamma = \tau(p)$, then pAp is isomorphic to $L(F_{1+(t-1)/(\gamma/\lambda)^2})$. Thus $\text{rdim}(pAp) = \gamma^2(1 + (t-1)/(\gamma/\lambda)^2 - 1) = \lambda^2(t - 1)$, as desired.

For a matrix algebra, we see compression does not change the size of the minimal projection (note it is important that $p \in A$), and thus does not change the regulated dimension. For diffuse hyperfinite algebras we see that the regulated will of course remain zero.

Since the regulated dimension is additive over direct sums, and since p has full central support (thus all factors in A are represented in pAp , the result follows. $\square$

Clearly this also means that the regulated dimension is invariant over dilatation.

Unfortunately unlike the free dimension, the regulated dimension can be negative and does not match the number of generators in an free group factor. It also may not be well defined in the same way as the free dimension if the interpolated free group factors are isomorphic.

Importantly if $\text{fdim}(A *_D B) = \text{fdim}(A) + \text{fdim}(B) - \text{fdim}(D)$ then $\text{rdim}(A *_D B) = \text{rdim}(A) + \text{rdim}(B) - \text{rdim}(D)$.

We can use the fact that this is invariant under dilation to extend the definition to $\mathcal{R}_4$. Since for any n , $\text{rdim}(L(F_s) \otimes M_n) = t^2(s-1)$ it is natural define $\text{rdim}(L(F_s) \otimes B(\mathcal{H})) = t^2(s-1)$. Note that $B(\mathcal{H}) \otimes L(F_s) \cong B(\mathcal{H}) \otimes L(F_{1+t^2(s-1)/t^2})$ with a trace preserving isomorphism, however note in both cases rdim is $t^2(s-1)$. While it is known that all factors of the form $B(\mathcal{H}) \otimes L(F_s)$ are isomorphic, there is only a trace preserving isomorphism if the interpolated free group factors are isomorphic.

Similarly for $B(\mathcal{H})$ we naturally define the regulated dimension to be $-t^2$, as it is for any n with M_n . Of course for any diffuse semifinite hyperfinite algebra we will set the regulated dimension to zero.

Using the above and the fact that the regulated dimension additive over direct sums, we can then define it for $\mathcal{R}_4$, except when this gives us $\infty - \infty$.

Definition 3.2. Let A be a semifinite von Neumann algebra in $\mathcal{R}_4$, written as:

$$A = H \oplus \bigoplus_{i \in I}^{p_i} L(F_{s_i}) \oplus \bigoplus_{j \in J} (L(F_{s_j}) \otimes B(\mathcal{H})) \oplus \bigoplus_{k \in K} M_{n_k} \oplus \bigoplus_{\ell \in L} B(\mathcal{H}),$$

where H is a diffuse hyperfinite algebra. Then

$$\text{rdim}(A) = \sum_{i \in I} \tau(p_i)^2 (s_i - 1) + \sum_{j \in J} t_j^2 (s_j - 1) - \sum_{k \in K} t_k^2 - \sum_{\ell \in L} t_\ell^2,$$

if this is defined.

Notation 3.3. To simplify things we use $\mathcal{F}_s^t$ to denote $L(F_{1+s/t^2})$ for $0 < t < \infty$ equipped with a trace so that $\tau(I_{\mathcal{F}_s^t}) = t$ and $B(\mathcal{H}) \otimes L(F_{1+s})$ if $t = \infty$. We will refer to these in general as semifinite interpolated free group factors.

Using this we can rewrite the definition of regulated dimension in a simpler way. For $A = \bigoplus_{i \in I} \mathcal{F}_{s_i}^{t_i} \oplus \bigoplus_{j \in J} M_{n_j}$, where n_j may be infinity, then $\text{rdim}(A) = \sum_{i \in I} s_i - \sum_{j \in J} t_j^2$.

Example 3.4. Consider the algebra in $\mathcal{R}_4$ as follows $\bigoplus_{i=1}^{\infty} (\mathcal{F}_1^1 \oplus \mathbb{C})$. If we compute the regulated dimension we see that $\text{rdim}(A) = \infty - \infty$ and thus is not well defined.

3.2. Substandard Embeddings. When working with semifinite algebras it makes more sense to extend the concept of the standard embedding, which we do in the following way:

Definition 3.5. We say that a trace preserving embedding $\phi : \mathcal{F}_s^t \rightarrow \mathcal{F}_{s'}^{t'}$ is *substandard* if for some projection $p \in \mathcal{F}_s^t$ with finite (non-zero) trace, the restriction $\phi|_{p\mathcal{F}_s^t p} : p\mathcal{F}_s^t p \rightarrow \phi(p)\mathcal{F}_{s'}^{t'}\phi(p)$ is a standard embedding or an isomorphism.

Note it is clearly equivalent to say this is true for any finite non-zero trace projection. Also note that substandard embeddings give us increasing regulated dimension, but not necessarily increasing free dimension.

The first three properties of the standard embedding listed before follow directly for the substandard embedding. The final is as follows:

Proposition 3.6. *For $\{\mathcal{F}_{s_i}^{t_i}\}_{i=1}^\infty$ and substandard embeddings $\phi_i : \mathcal{F}_{s_i}^{t_i} \rightarrow \mathcal{F}_{s_{i+1}}^{t_{i+1}}$ then the inductive limit is $\mathcal{F}_s^t$ where $s = \lim_{i \rightarrow \infty} s_i$ and $t = \lim_{i \rightarrow \infty} t_i$, and the induced embedding of $\mathcal{F}_{s_i}^{t_i} \rightarrow \mathcal{F}_s^t$ is substandard.*

Proof. Let $\rho_i = \phi_i \circ \phi_{i-1} \circ \cdots \circ \phi_1$. Choose a projection $p \in \mathcal{F}_{s_1}^{t_1}$ with finite non-zero trace t' .

Then $\rho_i|_p$ considered as map from $p\mathcal{F}_{s_1}^{t_1}p \rightarrow \rho_i(p)\mathcal{F}_{s_{i+1}}^{t_{i+1}}\rho_i(p)$ is standard or an isomorphism. Furthermore we know $\rho_i(p)\mathcal{F}_{s_{i+1}}^{t_{i+1}}\rho_i(p) \cong \mathcal{F}_{s_{i+1}}^{t'}$. Then we see each $\rho_{i-1}(p)\mathcal{F}_{s_i}^{t_i}\rho_{i-1}(p)$ is generated by R_i and S_i , where R_i is a copy of the hyperfinite II_1 factor, and $S_i = \{q_j X_j q_j\}_{j \in J_i}$ with $\sum_{j \in J_i} \tau(q_j)^2 = s_i$, where the X_j are a semi-circular system and $q_j \in R_i$, and so that $\phi_i(R_i) = R_{i+1}$ and $\phi_i(S_i) \subseteq S_{i+1}$. Thus $\rho_i(p)\mathcal{F}_{s_{i+1}}^{t_{i+1}}\rho_i(p) \cong \mathcal{F}_{s_{i+1}}^{t'}$.

Then for each i , $\mathcal{F}_{s_i}^{t_i}$ is generated by S_i and a dilation of R_i . The inductive limit is then generated by $S = \cup_{i=1}^\infty S_i$ and R , a hyperfinite II_1 (if $\lim_i t_i < \infty$) or II_∞ (if $\lim_i t_i = \infty$) factor. If we let $J = \cup_i J_i$ then we see $\sum_{j \in J} \tau(q_j)^2 = \lim_i s_i$. Thus the inductive limit is $\mathcal{F}_s^t$ as desired. $\square$

Finally, for convenience of notation we will further extend the definition of substandard in the following way:

Definition 3.7. For von Neumann algebras A and B in $\mathcal{R}_4$, an embedding $\phi : A \rightarrow B$ is said to be substandard if for every semifinite interpolated free group factor summand of A , $\mathcal{F}_s^t$ with central support p , $\phi : \mathcal{F}_s^t \rightarrow \phi(p)B\phi(p)$ is substandard.

3.3. Main Lemmas. The following lemma, based on Lemma 5.2 in [4] will be used for almost all of our results.

Lemma 3.8. *Let $\mathcal{R}$ be a class of von Neumann algebras such that if $A \in \mathcal{R}$ and $p \in A$ then $pAp \in \mathcal{R}$. Let A and B be von Neumann algebras in $\mathcal{R}$ with atomic type I subalgebra D . If for all A' and B' in $\mathcal{R}$ with abelian multimatrix subalgebra D' , $A' *_D B'$ is in $\mathcal{R}_4$, then so is $A *_D B$. Furthermore conditions on the number of $\mathcal{F}_s^t$ factor summands and type I atomic factor summands are preserved, and if we know $\text{rdim}(A' *_D B') = \text{rdim}(A') + \text{rdim}(B') - \text{rdim}(D')$ then the same holds for A , B , and D . Also if A and B are finite, then $A *_D B$ is too, and we can then make the same statement about $\mathcal{R}_3$.*

Proof. Let $D = \bigoplus_{k \in I_D} M_{n_k}^{p_k^D}$ (where we allow $n_k = \infty$, in which case we mean $B(\mathcal{H})$). For each $k \in I_D$, let $\{e_{i,j}^{(k)}\}$ be the standard basis for $M_{n_k}^{p_k^D}$. Let $e = \sum_{k \in I_D} e_{1,1}^{(k)}$ (with the sum converging in the WOT). Note this is a projection in D with central support I , and eDe is abelian. Let $A' = eAe$, $B' = eBe$, and $D' = eDe$. Let $\mathcal{M} = A *_D B$ and let $\mathcal{M}' = vN(A' \cup B')$, and since $A' \subseteq A$, $B' \subseteq B$ we see that A' and B' are free with amalgamation over D' , and thus $\mathcal{M}' = A' *_D B'$. Let $V = \{e_{1,j}^{(k)} | k \in I_D, 1 < j \leq n_k^D\}$. Note then that $A = vN(A' \cup V)$, $B = vN(B' \cup V)$, and $D = vN(D' \cup V)$. Then $\mathcal{M} = vN(\mathcal{M}' \cup V)$. Thus $\mathcal{M}' = e\mathcal{M}e$. By our assumption we know A' and B' are in $\mathcal{R}$ and D' is an abelian multimatrix algebra. Thus by assumption $\mathcal{M}'$ is of the correct form, which implies $\mathcal{M}$ is of the same form with the same number of factor summands of each kind (where the kinds are: diffuse hyperfinite, type I atomic, and $\mathcal{F}_s^t$ as this step may change interpolated free group factors to $\mathcal{F}_t^\infty$ or matrix algebras to $B(\mathcal{H})$). Now if $\text{rdim}(A' *_D B') = \text{rdim}(A') + \text{rdim}(B') - \text{rdim}(D')$, then since the central support of e is the identity $\text{rdim}(A *_D B) = \text{rdim}(A) + \text{rdim}(B) - \text{rdim}(D)$. $\square$

Lemma 3.9. *Let A be a semifinite hyperfinite von Neumann algebra with trace preserving conditional expectation onto a subalgebra $D = \bigoplus_{n \in N} \bigoplus_{t_n}^{p_n} \mathbb{C}$, for some countable index set N , so that the trace is semifinite on D as well. Then there exists a chain of multimatrix subalgebras A_j , so that $D \subseteq A_1 \subseteq A_2 \dots$ and $\cup_{j=1}^\infty A_j$ is dense in A .*

Proof. First we assume A is finite and begin by dividing it into type I and type II parts and approximating them separately. If A is type I and finite, and thus of the form $B \oplus (C \otimes L^\infty([0, 1]))$, where B and C are multimatrix algebras. We can choose a representation so that every p_n , $n \in N$, is composed of diagonal matrices in B and C and characteristic functions in $L^\infty([0, 1])$. Let $C = \bigoplus_{k \in K} C_k$ where each C_k is a matrix algebra. Then for each k in K we can choose a partition P_k of $[0, 1]$ into a countable number of measurable subsets so that for every $S \in P_k$ and $e_{j,j}^{(k)} \in C_k$, $e_{j,j}^{(k)} \otimes \chi_S \leq p_n$ for some $n \in N$. Then we set $A_1 = B \oplus \bigoplus_{k \in K} (C_k \otimes L^\infty(P_k))$, and note $D \subseteq A_1$. We can then define each A_j by further refining P_k to get the desired sequence.

For the finite type II case, we note that our algebra is isomorphic to $L^\infty(X, \mu) \otimes R$ for some measure μ . As before we pick subalgebras M_{2^k} of R so that $\cup_{k=1}^\infty M_{2^k}$ is dense in R , and denote the standard basis elements of M_{2^k} by $e_{i,j}^{(k)}$, and use the inclusion $M_{2^k} \rightarrow M_{2^{k+1}}$ where $e_{i,j}^{(k)} \rightarrow e_{2i-1, 2j-1}^{(k+1)} + e_{2i, 2j}^{(k+1)}$.

Let $f_n \in L^\infty(X, \mu)$ be the centre-valued trace of p_n in A , for all $n \in N$. Then let $S_{n,k} = f_n^{-1} \left(\bigcup_{\ell=1}^{2^{k-1}} \left[\frac{2\ell-1}{2^k}, \frac{2\ell}{2^k} \right) \right)$ for $k \geq 1$ and let $S_{n,0} = f_n^{-1}(\{1\})$. Then note $f_n = \sum_{k=0}^\infty \frac{1}{2^k} \chi_{S_{n,k}}$.

Now define $P_{0,k}$ to be a countable partition of X so that for every $n \in N$, $S_{n,k}$ with non-zero measure is the union of sets in $P_{0,k}$, and so that $P_{0,k}$ refines $P_{0,k-1}$. For each $S \in P_{0,k}$, let $s_{S,k}$ be the number of $n \in N$ such that $S \subseteq S_{n,k}$, and let $r_{S,k} = s_{S,k} + 2r_{S',k-1}$ where $S \subseteq S' \in P_{0,k-1}$, (for convenience we define $P_{0,-1} = \{X\}$ and $r_{X,-1} = 0$).

Then define A_0 to be the multimatrix algebra spanned by the basis

$$B_0 = \{e_{i,j}^{(k)} \otimes \chi_S | 0 \leq k, S \in P_{0,k}, r_{S',k-1} < i, j \leq r_{S,k}, S \subseteq S' \in P_{0,k-1}\}.$$

Note then for any $x \in X$ (except possibly on a set of measure zero),

$$\sum_{\substack{e_{i,i}^{(k)} \otimes \chi_S \in B_0, \\ x \in S}} e_{i,i}^{(k)} = I_R.$$

To check this, consider the trace of this sum. This is bounded above by $\sum_{n=1}^{\infty} f_n(x) = 1$, and for any $\epsilon > 0$ we can find an n_0 so that $\sum_{n=1}^{n_0} f_n(x) > 1 - \epsilon$ and a k_0 so that $2^{-k_0}n_0 < \epsilon$. Thus the sum of $\{\tau(e_{i,i}^{(k)}) = 2^{-k} | e_{i,i}^{(k)} \otimes \chi_S \in B_0, x \in S, k \leq k_0\}$ is at least $1 - 2\epsilon$. It follows from our choice of $r_{S,k}$ that the diagonal basis elements are orthogonal. It is also clear that this algebra contains a set of orthogonal projections p'_n with centre-valued traces f_n for $n \in N$. These are then equivalent to the p_n in D , and thus without loss of generality we can identify them with D .

To define B_m , we first define $P_{m,k}$ to be a refinement of $P_{m-1,k}$ such for any $S \in P_{m,k}$ either $0 < \mu(S) \leq 2^{-m}$ or S is a single atom. We define $r_{S,k}$ and $s_{S,k}$ in the same way (note for $S \in P_{m,k}$ if there exists an $S' \in P_{m',k}$ with $S \subseteq S'$ and $m > m'$ then $r_{S,k} = r_{S',k}$ and $s_{S,k} = s_{S',k}$ so this is well defined, and we do not need to add an index m).

Then let A_m be the multimatrix algebra spanned by

$$B_m = \{e_{i,j}^{(k)} \otimes \chi_S | m < k, S \in P_{m,k}, r_{S',k-1} < i, j \leq r_{S,k}, S \subseteq S' \in P_{m,k-1}, \\ \text{or if } k = m, S \in P_{m,k}, 1 \leq i, j \leq r_{S,m}\}.$$

Recalling the inclusion of $M_{2^{m-1}}$ into M_{2^m} and that each $P_{m,k}$ is a refinement of $P_{m-1,k}$, we see $A_{m-1} \subseteq A_m$.

We have established that for any $x \in X$ (except possibly on a set of measure zero) as m goes to infinity the sum of $\{\tau(e_{i,i}^{(k)}) = 2^{-k} | e_{i,i}^{(k)} \otimes \chi_S \in B_0, x \in S, k \leq m\}$ goes to 1. Since for A_m there is a set S containing x with measure less than $\max\{\mu(\{x\}), 2^{-m}\}$ and matrix subalgebra $M_{m'} \otimes \chi_S \subseteq M_{2^m} \otimes \chi_S$, whose trace is $\mu(S)$ times that sum. Thus this is dense in $\cup_m M_{2^m} \otimes L^\infty(X, \mu)$ and thus dense in A .

If A is only semifinite, then let q_k be the sum of the first k p_n in D . Then use the finite case to construct an approximating chain $A_{i,1}$ of $q_1 A q_1$ which contains $q_1 D q_1$. Then use the finite case again to let $A_{i,k}$ be an approximating chain of $q_k A q_k$ containing $q_k D q_k$ and so that $A_{i,k} \subseteq A_{i,k+1}$ for all i and for $k \geq 1$ (note

the construction above allows this). Then define our approximating chain to be $A_i = A_{i,i} \oplus (1 - q_i)D(1 - q_i)$.

□

Definition 3.10. We call an embedding $\phi : \mathcal{N} \rightarrow \mathcal{M}$ a *simple step* if it follows one of the two following patterns:

$$(1) \mathcal{N} = \overset{p}{A} \oplus \overset{q}{B}, \mathcal{M} = \left(\bigoplus_{i=1}^n \overset{p_i}{A} \right) \oplus \overset{q}{B}, \text{ with } p = \sum_{i=1}^n p_i, \text{ and } \phi(a, b) = (a, \dots, a, b).$$

We call this a *simple step of the first kind*.

$$(2) \mathcal{N} = M_n \oplus_t M_m \oplus \overset{p}{B}, \mathcal{M} = M_{n+m} \oplus \overset{p}{B} \text{ where } \phi(x, y, b) = \left(\begin{bmatrix} x & 0 \\ 0 & y \end{bmatrix}, b \right). \text{ We call this a simple step of the second kind.}$$

Lemma 3.11. For two multimatrix algebras $\mathcal{N}$ and $\mathcal{M}$ and a trace preserving inclusion $\phi : \mathcal{N} \rightarrow \mathcal{M}$, then ϕ can be written as a composition of a (possibly countably infinite) sequence of simple steps.

Proof. The proof is identical to Lemma 3.5 in [6], following directly from examining either the Bratteli diagrams or the minimal projections. The only difference is that since we have an infinite number of matrix algebras, each of which can be split into an infinite number of matrix algebras, we may need an infinite number of steps. □

3.4. Main Theorem and Examples.

Theorem 3.12. Let A and B hyperfinite von Neumann algebras with trace preserving conditional expectations onto a type I atomic subalgebra D then:

- (1) If A and B are finite then $A *_D B$ is in $\mathcal{R}_3$
- (2) If A and B are semifinite, and the trace on D is still semifinite, then $A *_D B$ is in $\mathcal{R}_4$.

Furthermore $\text{rdim}(A *_D B) = \text{rdim}(A) + \text{rdim}(B) - \text{rdim}(D)$ if this is well defined.

Proof. By Lemma 3.8 assume without loss of generality that D is abelian. If D is finite dimensional we can apply Theorem 2.9 (since A and B aren't finite, then finite dimensional D is not semifinite), so we may also assume $D = \bigoplus_{k=1}^{\infty} \overset{p_k^D}{t_k^D} \mathbb{C}$.

Use Lemma 3.9 to define the sequences A_i and B_j to be chains of subalgebras of A in B , respectively, containing D . Note in the construction from Lemma 3.9, the atomic parts of A and B are included in every A_i and B_j . Let $q_k^D = \sum_{k'=1}^k p_{k'}^D$. Then let

$$\mathcal{M}(i, j, k) = \left(q_k^D A_j q_k^D *_{{q_k^D D q_k^D}} q_k^D B_j q_k^D \right),$$

and let $\mathcal{N}(i, j, k)$ be

$$\left(q_{k-1}^D A_j q_{k-1}^D *_{{q_{k-1}^D D q_{k-1}^D}} q_{k-1}^D B_j q_{k-1}^D \right) \oplus \left(p_k^D A_i p_k^D *_{{p_k^D D p_k^D}} p_k^D B_j p_k^D \right).$$

For fixed k , the sequence $\mathcal{M}(i, j, k)$, is an approximating sequence for the amalgamated free product of hyperfinite von Neumann algebras over $q_k^D D$ (which is finite dimensional), as in Theorem 2.9. Thus we can choose i_k and j_k sufficiently large to be the start of the sequence in the proof of that theorem. In particular, for any minimal projection $p \in A_{i_k}$ (resp B_{j_k}) from the diffuse part of A (resp. B), so that $p \leq p_{k'}^D$ with $k' \leq k$ then $\tau(p) < t_{k'}^D - \tau(p')$ for any minimal projection $p' \in B$ (resp A) with $p' \leq p_{k'}^D$, unless $p_{k'}^D$ is minimal in B (resp. A), and also so that the number of interpolated free group factors in $\mathcal{M}(i, j, k)$ is stable for all $i \geq i_k$ and $j \geq j_k$. Furthermore we can assume that if $\text{rdim}(A)$ (resp B) is finite then $\text{rdim}(q_k A_{i_k} q_k) > \text{rdim}(q_k A q_k) - 1/k$.

Our goal is to show that the inclusions of the sequence

$$\mathcal{M}(i_2, j_2, 1) \rightarrow \mathcal{N}(i_2, j_2, 2) \rightarrow \mathcal{M}(i_2, j_2, 2) \rightarrow \mathcal{M}(i_3, j_3, 2) \rightarrow \mathcal{N}(i_3, j_3, 3) \rightarrow \dots$$

are substandard.

First examine the steps of the form $\mathcal{M}(i_k, j_k, k) \rightarrow \mathcal{M}(i_{k+1}, j_{k+1}, k)$. Since k does not change in this step, it is just like the steps in the proof of Theorem 2.9 (from [6]). Using Lemma 3.11, we break this up into a sequence of inclusions induced by simple steps in the subalgebras of A or B , so we need only show that this type of inclusion is substandard. Consider the inclusion $\mathcal{M}(i, j, k) \rightarrow \mathcal{M}(i+1, j, k)$ with $i_k \leq i < i_{k+1}$ and $j_k \leq j \leq j_{k+1}$ where $A_i \rightarrow A_{i+1}$ is a simple step (note this may not work precisely with our original indexing, but this does not affect the proof and eases notation).

If the inclusion $A_i \rightarrow A_{i+1}$ is a simple step of the first kind, which makes copies of a summand, M_n , of A_i , then either M_n is contained in some interpolated free group factor summand, or it is the corner of some matrix factor summand (as in the proof of Theorem 2.9). In the former case we apply Lemma 2.8 to show this is a standard embedding. In the latter we see the matrix factor is similarly copied.

Alternately, if the inclusion $A_i \rightarrow A_{i+1}$ is a simple step of the second kind, then we are adding a partial isometry v connecting two matrix algebras in A_i . Again we proceed exactly as in Theorem 2.9. If both vv^* and v^*v are in an interpolated free group factor we apply Lemma 2.7 to show this is a standard embedding. If only one of vv^* and v^*v is in an interpolated free group factor we apply Lemma 2.5 to show that this is a substandard embedding. If neither vv^* nor v^*v are in interpolated free group factors, then they are both in matrix algebras, then these are embedded in another matrix algebra.

Thus applying Proposition 3.6 to this sequence of inclusions we see then that any interpolated free group factor summand in $\mathcal{M}(i_{k-1}, j_{k-1}, k)$, is embedded in another in $\mathcal{M}(i_k, j_k, k)$ by a substandard embedding, and so this inclusion is substandard.

Next we examine the inclusions of the form $\mathcal{N}(i_k, j_k, k) \rightarrow \mathcal{M}(i_k, j_k, k)$. This is similar to the proof of Theorem 2.6 (from [4]), since i_k and j_k do not change, and thus we have a fixed pair of multimatrix algebras A_{i_k} and B_{j_k} .

Let $q_k A_{i_k} q_k = \bigoplus_{\ell \in I_{A_{i_k}}} M_{n_\ell}^{p_\ell}$ and $q_k B_{j_k} q_k = \bigoplus_{\ell \in I_{B_{j_k}}} M_{n_\ell}^{p_\ell}$. Now for each $\ell \in I_{A_{i_k}} \cup I_{B_{j_k}}$, if p_ℓ is not orthogonal to either q_{k-1} or p_k , then choose a $v_\ell \in M_{n_\ell}$ which is a partial isometry so that $v_\ell v_\ell^* \leq p_k$ and $v_\ell^* v_\ell \leq q_{k-1}$, and each is minimal in A_{i_k} of B_{j_k} . Let V be the set of these v_ℓ . Then note that $\mathcal{N}(i_k, j_k, k) \cup V$ generates $\mathcal{M}(i_k, j_k, k)$. Let $\mathcal{N}(i_k, j_k, k, V') = vN(\mathcal{N}(i_k, j_k, k) \cup V')$ for $V' \subseteq V$. We show that for $v \in V$ and $V' \subseteq V \setminus \{v\}$, the inclusion $\mathcal{N}(i_k, j_k, k, V') \rightarrow \mathcal{N}(i_k, j_k, k, V' \cup \{v\})$ is substandard.

Let $q = vv^* + v^*v$, and examine $q\mathcal{N}(i_k, j_k, k, V')q$, and the algebra M_2 generated by v . As in the proofs of Theorem 2.6 and Theorem 2.9, we see that

$$q\mathcal{N}(i_k, j_k, k, V')q *_{\substack{\mathbb{C} \\ \oplus \\ \mathbb{C}}}^{vv^* \quad v^*v} M_2 = q\mathcal{N}(i_k, j_k, k, V' \cup \{v\})q.$$

If either of vv^* or v^*v are minimal and central in $q\mathcal{N}(i_k, j_k, k, V')q$, (without loss of generality assume vv^*), then vv^* is a minimal projection in a matrix algebra summand M_n in $\mathcal{N}(i_k, j_k, k, V')$. Any matrix algebra summand $M_{m'}$ under v^*v in $q\mathcal{N}(i_k, j_k, k, V')q$ must be part of a matrix algebra summand M_m in $\mathcal{N}(i_k, j_k, k, V')$. Then the inclusion of M_m in $\mathcal{N}(i_k, j_k, k, V' \cup \{v\})$ maps it to the corner of a matrix summand $M_{m+nm'}$. Any interpolated free group factor under v^*v , is included into a dilation of itself, which is a substandard embedding. Finally if there is a diffuse hyperfinite algebra under v^*v then it is included in a diffuse hyperfinite algebra.

If neither vv^* nor v^*v is minimal and central, then we can apply Lemma 2.5. This shows us that the inclusion of $q\mathcal{N}(i_k, j_k, k, V')q$ into $q\mathcal{N}(i_k, j_k, k, V' \cup \{v\})q$ is substandard, and thus so is the inclusion of $\mathcal{N}(i_k, j_k, k, V')$ into $\mathcal{N}(i_k, j_k, k, V' \cup \{v\})$. Thus, applying Proposition 3.6, so is the inclusion $\mathcal{N}(i_k, j_k, k) \rightarrow \mathcal{M}(i_k, j_k, k)$.

In the final case, $\mathcal{M}(i_k, j_k, k-1) \rightarrow \mathcal{N}(i_k, j_k, k)$ is immediate.

Thus the inclusions in the chain,

$$\mathcal{M}(i_2, j_2, 1) \rightarrow \mathcal{N}(i_2, j_2, 2) \rightarrow \mathcal{M}(i_2, j_2, 2) \rightarrow \mathcal{M}(i_3, j_3, 2) \rightarrow \mathcal{N}(i_3, j_3, 3) \rightarrow \dots,$$

are all substandard. At all stages this is the direct sum of a finite number of interpolated free group factors and a hyperfinite algebra. Thus, applying Proposition 3.6 once more, the inductive limit is a countable direct sum of factors of the form $\mathcal{F}_s^t$ and a hyperfinite algebra, and thus in $\mathcal{R}_4$. If A and B were finite, then clearly $A *_D B$ is finite, and thus in $\mathcal{R}_3$.

If all the regulated dimensions are defined, as well as the sum then it is clear that $\text{rdim}(A *_D B) = \lim_{k \rightarrow \infty} \text{rdim}(\mathcal{M}(i_k, j_k, k))$. Choose $k \in I_D$ and denote $A_{i_k} =$

$$\bigoplus_{\ell \in I_A} M_{n_\ell}^{p_\ell}_{t_\ell}.$$
 Then

$$\text{rdim}(q_k A_{i_k} q_k) = - \sum_{\ell \in I_A, p_\ell q_k \neq 0} t_\ell^2$$

and

$$\text{rdim}(A_{i_k}) = - \sum_{\ell \in I_A} t_\ell^2.$$

For every $\ell \in I_D$ so that $p_\ell q_k = 0$, there is a minimal projection $p'_\ell \in M_{n_\ell}$ less than some $p_{k'}^D$ with $k' > k$. This tells us that

$$\sum_{\ell \in I_D, p_\ell q_k = 0} t_\ell^2 < \sum_{k' > k} (t_{k'}^D)^2.$$

Thus

$$|\text{rdim}(q_k A_{i_k} q_k) - \text{rdim}(A_{i_k})| < \left| \sum_{k' > k} (t_{k'}^D)^2 \right| = \text{rdim}((1 - q_k)D(1 - q_k)).$$

Thus if $\text{rdim}(D) \neq -\infty$ then $\lim_{k \rightarrow \infty} \text{rdim}(q_k A_{i_k} q_k) = \text{rdim}(A)$. The same calculation works for B_{j_k} . Then note by Theorem 2.9

$$\text{rdim}(\mathcal{M}(i_k, j_k, k)) = \text{rdim}(q_k A_{i_k} q_k) + \text{rdim}(q_k B_{j_k} q_k) - \text{rdim}(q_k D q_k).$$

Then if $\text{rdim}(D) \neq -\infty$ then

$$\begin{aligned} \lim_{k \rightarrow \infty} \text{rdim}(\mathcal{M}(i_k, j_k, k)) &= \lim_{k \rightarrow \infty} \text{rdim}(A_{i_k}) + \lim_{k \rightarrow \infty} \text{rdim}(B_{j_k}) - \text{rdim}(D) \\ &= \text{rdim}(A) + \text{rdim}(B) - \text{rdim}(D). \end{aligned}$$

If $\text{rdim}(D) = -\infty$ then we see that if $\text{rdim}(A)$ and $\text{rdim}(B)$ are not $-\infty$, so $\text{rdim}(q_k A_{i_k} q_k) + \text{rdim}(q_k B_{j_k} q_k)$ is bounded below, while $-\text{rdim}(q_k D q_k)$ goes to ∞ , the equation again holds.

□

Example 3.13. Let:

$$\begin{aligned} D &= \bigoplus_{i=1}^{\infty} \overset{p_i^D}{\mathbb{C}}_{\frac{1}{i}} \\ A &= \bigoplus_{i=1}^{\infty} \overset{p_i^A}{R}, \tau(p_i^A) = \frac{1}{2i-1} + \frac{1}{2i} \\ B &= \overset{p_0^B}{R} \oplus \bigoplus_{i=1}^{\infty} \overset{p_i^B}{R}, \tau(p_i^B) = \frac{1}{2i} + \frac{1}{2i+1}, \tau(p_0^B) = 1. \end{aligned}$$

Here we set $p_{2i-1}^D + p_{2i}^D = p_i^B$ and $p_{2i}^D + p_{2i+1}^D = p_i^B$, with $p_0^B = p_1^D$. We can see that $r = \text{rdim}(A) + \text{rdim}(B) - \text{rdim}(D) = \frac{\pi^2}{6}$. From there it is not hard to check that $A *_D B = \mathcal{F}_{\frac{\pi^2}{6}}^\infty$. This shows that even if all our factor summands are finite, the result may have a non-finite factor summand.

Example 3.14. Let

$$D = \bigoplus_{i=1}^{\infty} \mathbb{C}_1$$

$$A = B = \bigoplus_{i=1}^{\infty} \mathbb{C}_{1/2}.$$

Then computing we see

$$\text{rdim}(A) + \text{rdim}(B) - \text{rdim}(D) = -\infty - \infty + \infty,$$

which is not defined. Computing the product we see that $A *_D B = \bigoplus_{i=1}^{\infty} L(\mathbb{Z}) \otimes M_2$. In this case $\text{rdim}(A *_D B) = 0$, however it would be easy to add a summand so that we could get any value from this.

4. CLOSEDNESS OF $\mathcal{R}_3$ AND $\mathcal{R}_4$

The following lemma was proved as Theorem 3.1 in [1].

Lemma 4.1. *Let R and R' be hyperfinite II_1 factors. Let $D = \bigoplus_{i \in I} \mathbb{C}_{t_i}^{p_i}$ (I countable) be a common subalgebra of R and R' . Then $R *_D R' = L(F_{1+\sum_{i \in I} t_i^2})$, and this is generated by $R' \cup \{p_i X_i p_i\}_{i \in I}$.*

Proof. The fact that $R *_D R'$ is the interpolated free group factor mentioned follows from Theorem 3.12, so here we need only check that it is generated correctly.

First assume that I is finite, and proceed by induction on the size of I . In the case $|I| = 1$, this has been established in Corollary 3.6 in [3]. Let i_0 be such that t_{i_0} is minimal. Using Theorem 2.1 (b) from [5], we define

$$\begin{aligned} \mathcal{N}_1 &= vN((1 - p_{i_0})R(1 - p_{i_0}) \cup (1 - p_{i_0})R'(1 - p_{i_0})) \\ &= (1 - p_{i_0})R(1 - p_{i_0}) *_{(1-p_{i_0})D} (1 - p_{i_0})R'(1 - p_{i_0}). \end{aligned}$$

Then by our induction hypothesis, $\mathcal{N}_1 = L(F_{1+\sum_{i \in I, i \neq i_0} \frac{t_i^2}{(1-t_{i_0})^2}})$, generated by $(1 - p_{i_0})R'(1 - p_{i_0}) \cup \{p_i X_i p_i\}_{i \in I \setminus \{i_0\}}$

Then using part (c) of Theorem 2.1 in [5], we see that $r = p_{i_0}$, thus $\tilde{A} = (1 - p_{i_0})R(1 - p_{i_0}) \oplus \mathbb{C}p_{i_0}$, and thus $p_{i_0}\tilde{A}p_{i_0} = \mathbb{C}$. From this we see directly that for $\mathcal{N}_2 = vN(\tilde{A} \cup R')$, $\mathcal{N}_1 = (1 - p_{i_0})\mathcal{N}_2(1 - p_{i_0})$. Thus we write $\mathcal{N}_2 = vN(R' \cup \{p_i X_i p_i\}_{i \in I, i \neq i_0})$.

Using part (d) of Theorem 2.1 in [5], we see that since $p_{i_0}(R *_D R')p_{i_0} = p_{i_0}\mathcal{N}_2p_{i_0} * L(\mathbb{Z})$, and that $R *_D R'$ is generated by $R' \cup \{p_i X_i p_i\}_{i \in I}$, and thus we are done.

For the infinite case, let $K = \mathbb{N}$, ordered so that $t_i \leq t_{i-1}$, and let $q_k = \sum_{i=1}^k p_i^D$. Let $\mathcal{M}_k = q_k R q_k *_{q_k D} q_k R' q_k$. The above tells us that each $\mathcal{M}_k$ is generated by $q_k R' q_k \cup \{p_i X_i p_i\}_{i=1}^k$ and that the embeddings of $\mathcal{M}_k \rightarrow \mathcal{M}_{k+1}$ are substandard, thus the generating set in the inductive limit $\mathcal{M}$ is as desired, completing the proof. $\square$

Corollary 4.2. *Let A and B each be copies of the hyperfinite II_∞ factor, with subalgebra $D = \bigoplus_{i=1}^{\infty} \mathbb{C}^{p_i}_{t_i}$ and trace preserving conditional expectation onto D so the trace is semifinite on D . Then $A *_D B = \mathcal{F}^{\infty}_{-rdim(D)}$, and this is generated by $A \cup \{p_i X_i p_i\}_{i=1}^{\infty}$*

Proof. Let $q_k = \sum_{i=1}^k p_i$. Then by Lemma 4.1 any $q_k A q_k *_D q_k B q_k$ is generated by $q_k A q_k \cup \{p_i X_i p_i\}_{i=1}^k$, thus the inductive limit is generated by the desired set. $\square$

Lemma 4.3. *Let $A = \mathcal{F}^t_{s_A}$ and $B = \mathcal{F}^t_{s_B}$, each with subalgebra $D = \bigoplus_{k \in K} \mathbb{C}^{p_k^D}_{t_k}$. Then $\mathcal{M} = A *_D B \cong \mathcal{F}^t_{s_A + s_B - rdim(D)}$, and the inclusion of $A \rightarrow \mathcal{M}$ is substandard.*

Proof. Let p be a minimal projection in D , and choose representations of A and B of the form $A = vN(R \cup \{p_i X_i p_i\}_{i \in I})$ and $B = vN(R' \cup \{q_j X_j q_j\}_{j \in J})$, where $p_i \leq p$ and $q_j \leq p$ for all $i \in I, j \in J$, and R and R' are either the hyperfinite II_1 factor or the hyperfinite II_∞ factor. Then $A *_D B$ is generated by $R \cup R' \cup \{p_i X_i p_i\}_{i \in I} \cup \{q_j X_j q_j\}_{j \in J}$. From Theorem 4.1 we know that $R *_D R' = \mathcal{F}^t_{-fdim(D)}$, with R embedded correctly and so $R *_D R'$ is generated by $R \cup \{p_k^D X_k p_k^D\}_{k \in K}$. We can then find $q'_j \in R$ which are unitary conjugates of q_j using unitaries in $R *_D R'$, thus $\mathcal{M}$ is generated by $R \cup \{p_k^D X_k p_k^D\}_{k \in K} \cup \{p_i X_i p_i\}_{i \in I} \cup \{q'_j X_j q'_j\}_{j \in J}$. Thus $\mathcal{M}$ is the desired factor, and the inclusion of A is substandard. $\square$

Lemma 4.4. *Let H be a hyperfinite von Neumann algebra and $A = \mathcal{F}^t_s$, with common abelian type I atomic subalgebra D , then $A *_D H = \mathcal{F}^t_{s+rdim(H)-rdim(D)}$, and the inclusion of $\mathcal{F}^t_s$ into it is substandard.*

Proof. Using Lemma 3.9 we can find a chain of multimatrix algebras H_i approximating H so that $D \subseteq H_i$ for all i , and that so that for each i there exists a finite projection $q_i \in D$ so that $(1 - q_i)H_i(1 - q_i) = (1 - q_i)D$. Applying Lemma 3.11 we can assume that this is generated by a sequence of simple steps. Then by Lemma 5.9 of [4], since $q_i A q_i$ is an interpolated free group factor, $q_i H_i q_i *_D q_i A q_i$ is an interpolated free group factor (with regulated dimension as expected), and the inclusion of $q_i A q_i$ is standard. Then apply Lemma 2.4 to see this implies $H_i *_D A$ is $\mathcal{F}^t_{s+rdim(H_i)-rdim(D)}$, and the inclusion of A into it is substandard.

Using Lemma 2.8, and Lemma 2.7 we see that the embedding of $H_i *_D A \rightarrow H_{i+1} *_D A$ is substandard. which completes the proof. $\square$

Lemma 4.5. *Let $A \in \mathcal{R}_4$ and $B = \mathcal{F}^t_s$. Let D be an abelian type I atomic subalgebra of both A and B . Then $\mathcal{M} = A *_D B$ is a semifinite interpolated free group factor, and if $rdim(A)$ is defined and greater than $-\infty$, then $rdim(\mathcal{M}) = \mathcal{F}^t_{s+rdim(A)-rdim(D)}$. Furthermore the inclusion $B \rightarrow \mathcal{M}$ is substandard and that of $F_i \rightarrow \mathcal{M}$ is substandard for any summand, $F_i \cong \mathcal{F}^{t'}_{s'}$, of A .*

Proof. Let $A = H \oplus \bigoplus_{i=1}^K \mathcal{F}_{s_i}^{t_i}$ for $K \in \mathbb{N} \cup \{\infty\}$ and where H is a hyperfinite algebra.

Let $A_0 = H \oplus \bigoplus_{i=1}^K p_i D$ and let $A_k = H \oplus \bigoplus_{i=1}^k \mathcal{F}_{s_i}^{t_i} \oplus \bigoplus_{i=k+1}^K p_i D$. Let $\mathcal{M}_k = A_k *_D B$.

Our goal is to show that $\mathcal{M}_k = \mathcal{F}_{\text{rdim}(A_k)+s-\text{rdim}(D)}^t$ for all k , and that the inclusion of $\mathcal{M}_k \rightarrow \mathcal{M}_{k+1}$ is substandard.

First Lemma 4.4 shows that $\mathcal{M}_0 = \mathcal{F}_{\text{rdim}(A_0)+s-\text{rdim}(D)}^t$, and the inclusion of $B \rightarrow \mathcal{M}_0$ is standard.

Lemma 2.4 tells us that $p_k \mathcal{M}_k p_k = p_k \mathcal{M}_{k-1} p_k *_D \mathcal{F}_{s_k}^{t_k}$ and that $\mathcal{M}_k$ is a factor. Thus the embedding of $\mathcal{F}_{s_k}^{t_k} \rightarrow p_k \mathcal{M}_k p_k$ is substandard. By Lemma 4.3, $p_k \mathcal{M}_k p_k = \mathcal{F}_{s'}^{t_k}$ for some s' and the inclusion of $p_k \mathcal{M}_{k-1} p_k$ into it is substandard, and thus $\mathcal{M}_k = \mathcal{F}_{\text{rdim}(A_k)+s-\text{rdim}(D)}^t$ and the inclusion $\mathcal{M}_{k-1} \rightarrow \mathcal{M}_k$ is substandard.

Thus the inductive limit of the $\mathcal{M}_k$, $A *_D B = \mathcal{F}_{\text{rdim}(A)+s-\text{rdim}(D)}^t$, the inclusion of B into it is standard, and the inclusions of $\mathcal{F}_{s_i}^{t_i} \rightarrow \mathcal{M}$ are substandard. $\square$

Theorem 4.6. *For $A, B \in \mathcal{R}_4$ (or $\mathcal{R}_3$) and D an atomic type I subalgebra of them, $A *_D B \in \mathcal{R}_4$ (or $\mathcal{R}_3$), and $\text{rdim}(A *_D B) = \text{rdim}(A) + \text{rdim}(B) - \text{rdim}(D)$ if this equation is well defined.*

Proof. As usual we use Lemma 3.8 to assume without loss of generality that D is

abelian. Let $A = H_A \oplus \bigoplus_{i=1}^{K_A} \mathcal{F}_{s_i}^{t_i}$ where $K_A \in \mathbb{N} \cup \{\infty\}$ similarly let $B = H_B \oplus \bigoplus_{j=1}^{K_B} \mathcal{F}_{s'_j}^{t'_j}$.

Then define $A_k = H_A \oplus \bigoplus_{i=1}^k \mathcal{F}_{s_i}^{t_i} \oplus \bigoplus_{i=k+1}^{K_A} p_i D$ and define B_k similarly.

Let $\mathcal{M}(i, j) = A_i *_D B_j$, then the inductive limit of the $\mathcal{M}(i, j)$ is $A *_D B$. We claim that each $\mathcal{M}(i, j)$ is in $\mathcal{R}_3$, and that the embedding of $\mathcal{M}(i, j)$ into $\mathcal{M}(i+1, j)$ is substandard. We proceed by induction on i, j .

Theorem 3.12 tells us that $\mathcal{M}(0, 0)$ is in $\mathcal{R}_4$, and that $\text{rdim}(\mathcal{M}(0, 0)) = \text{rdim}(A_0) + \text{rdim}(B_0) - \text{rdim}(D)$. By Lemma 2.4, $p_i \mathcal{M}(i, j) p_i = vN(p_i \mathcal{M}(i-1, j) p_i \cup \mathcal{F}_{s_i}^{t_i}) = p_i \mathcal{M}(i-1, j) p_i *_D \mathcal{F}_{s_i}^{t_i}$. Lemma 4.5 tells us that this is of the form $\mathcal{F}_s^t$ for some s and t and that the inclusion of any $\mathcal{F}_s^t$ factor summand in $p_i \mathcal{M}(i-1, j) p_i \rightarrow p_i \mathcal{M}(i, j) p_i$ is substandard. This also tells us that the inclusion of $\mathcal{F}_{s_i}^{t_i} \rightarrow p_i \mathcal{M}(i, j) p_i$ is substandard.

Since the central support of p_i in $\mathcal{M}(i-1, j)$ and $\mathcal{M}(i, j)$ is the same (again by Lemma 2.4), any $\mathcal{F}_s^t$ factor summand of $\mathcal{M}(i-1, j)$ which is orthogonal to p_i is identical in $\mathcal{M}(i, j)$. Thus the inclusion of $\mathcal{M}(i-1, j) \rightarrow \mathcal{M}(i, j)$ is substandard, as is the inclusion of $\mathcal{F}_{s_i}^{t_i} \rightarrow \mathcal{M}(i-1, j)$.

Next note that $\text{rdim}(q \mathcal{M}(i, j) q) = \text{rdim}(p_i \mathcal{M}(i, j) p_i)$ by Lemma 3.1. Note

$$\text{rdim}(p_i \mathcal{M}(i, j) p_i) = \text{rdim}(p_i \mathcal{M}(i-1, j) p_i) + \text{fdim}(\mathcal{F}_{s_i}^{t_i}) - \text{rdim}(p_i D).$$

Since q is the central support of p_i in both $\mathcal{M}(i, j)$ and $\mathcal{M}(i-1, j)$ we know

$$\text{rdim}(p_i \mathcal{M}(i-1, j) p_i) = \text{rdim}(q \mathcal{M}(i-1, j) q).$$

Thus we see

$$\text{rdim}(q\mathcal{M}(i, j)q) = \text{rdim}(q\mathcal{M}(i-1, j)q) + \text{rdim}(\mathcal{F}_{s_i}^{t_i}) - \text{rdim}(p_i D).$$

And since $(1-q)\mathcal{M}(i, j)(1-q) = (1-q)\mathcal{M}(i-1, j)(1-q)$, we see

$$\text{rdim}(\mathcal{M}(i, j)) = \text{rdim}(\mathcal{M}(i-1, j)) + \text{rdim}(\mathcal{F}_{s_i}^{t_i}) - \text{rdim}(p_i D).$$

Note $\text{fdim}(p_i A_i p_i) = \text{rdim}(\mathcal{F}_{s_i}^{t_i})$, and $\text{rdim}(p_i A_{i-1} p_i) = \text{rdim}(p_i D)$. Thus

$$\text{rdim}(A_i) - \text{rdim}(A_{i-1}) = \text{rdim}(\mathcal{F}_{s_i}^{t_i}) - \text{rdim}(p_i D).$$

Combining these we get:

$$\begin{aligned} \text{rdim}(\mathcal{M}(i, j)) &= \text{rdim}(A_{i-1}) + \text{rdim}(B_j) - \text{rdim}(D) + \text{rdim}(\mathcal{F}_{s_i}^{t_i}) - \text{rdim}(p_i D) \\ &= \text{rdim}(A_i) + \text{rdim}(B_j) - \text{rdim}(D). \end{aligned}$$

Thus our claim is proved, and the result follows. Note that since if A and B are finite, their amalgamated free product is, and thus if they were both in $\mathcal{R}_3$ then so will their amalgamated free product. $\square$

Example 4.7. Let $A = \bigoplus_{i=1}^{\infty} \mathcal{F}_{2^i}^{2^{-i}}$ where $\tau(p_i) = \frac{1}{2^i}$ and let $B = \mathcal{F}_2^1$, and choose any multimatrix subalgebra D of these. Checking, we see $\text{rdim}(A) = \sum_{i=1}^{\infty} 1 = \infty$. By Lemma 4.5 we know $A *_D B$ is an interpolated free group factor, and applying the free dimension formula we see that it must be $\mathcal{F}_{\infty}^1$.

This shows that unlike in the case of $\mathcal{R}_1$ and $\mathcal{R}_2$, restricting the interpolated free group factor summands in A and B to having only finite free dimension does not guarantee this for the product. We could restrict $\mathcal{R}_3$ to those with finite regulated dimension (or equivalently free dimension) and it would be still closed under these amalgamated free products. Similarly we could restrict $\mathcal{R}_4$ to those with finite defined regulated dimension, which would then be closed over amalgamated free products of type I atomic algebras with finite defined regulated dimension.

REFERENCES

- [1] Nathaniel P. Brown and Kenneth J. Dykema. Popa algebras in free group factors. *J. Reine Angew. Math.*, 573:157–180, 2004.
- [2] Ken Dykema. Free products of hyperfinite von Neumann algebras and free dimension. *Duke Math. J.*, 69(1):97–119, 1993.
- [3] Ken Dykema. Interpolated free group factors. *Pacific J. Math.*, 163(1):123–135, 1994.
- [4] Ken Dykema. Amalgamated free products of multi-matrix algebras and a construction of subfactors of a free group factor. *Amer. J. Math.*, 117(6):1555–1602, 1995.
- [5] Ken Dykema. A description of amalgamated free products of finite von Neumann algebras over finite-dimensional subalgebras. *Bull. Lond. Math. Soc.*, 43(1):63–74, 2011.
- [6] Ken Dykema and Daniel Redelmeier. The amalgamated free product of hyperfinite von Neumann algebras over finite dimensional subalgebras, 2011, arXiv:1110.5597.

- [7] A. Guionnet, V. Jones, and D. Shlyakhtenko. A semi-finite algebra associated to a subfactor planar algebra. *J. Funct. Anal.*, 261(5):1345–1360, 2011.
- [8] Vijay Kodiyalam and V. S. Sunder. On the Guionnet-Jones-Shlyakhtenko construction for graphs. *J. Funct. Anal.*, 260(9):2635–2673, 2011.
- [9] Sorin Popa. Markov traces on universal Jones algebras and subfactors of finite index. *Invent. Math.*, 111(2):375–405, 1993.
- [10] Dan Voiculescu. Symmetries of some reduced free product C^* -algebras. In *Operator algebras and their connections with topology and ergodic theory (Buşteni, 1983)*, volume 1132 of *Lecture Notes in Math.*, pages 556–588. Springer, Berlin, 1985.
- [11] Dan Voiculescu. Operations on certain non-commutative operator-valued random variables. *Astérisque*, (232):243–275, 1995. Recent advances in operator algebras (Orléans, 1992).

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA

E-mail address: delredel@alumni.uwaterloo.ca